

US EPA ARCHIVE DOCUMENT

County Distribution of Federally-listed Endangered, Threatened, Proposed and Candidate Species

Revised February 2014

County	Species	Status	Habitat
Adams	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. Roosts and forages in upland forests and woods.
	Kirtland's warbler <i>Setophaga kirtlandii</i>	Endangered	Young jack pine stands (5 to 25 years old)
	Whooping crane <i>Grus americanus</i>	**Non-essential experimental population	Open wetlands and lakeshores Whooping cranes have nested in this county
	Karner blue butterfly <i>Lycaeides melissa samuelis</i>	Endangered	Prairie, oak savanna, and jack pine areas with wild lupine
Ashland	Canada lynx <i>Lynx canadensis</i>	Threatened	While no resident populations are known from Wisconsin, the species occasionally occurs in northern forested areas, and counties listed are those with the highest likelihood of occurrence.
	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Piping plover <i>Charadrius melodus</i>	Endangered	Sandy beaches; bare alluvial and dredge spoil islands
	Piping plover <i>Charadrius melodus</i>	Critical Habitat Designated	
	Rufa red knot <i>(Calidris canutus rufa)</i>	Proposed Threatened	Along Lake Superior
Barron	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
Bayfield	Canada lynx <i>Lynx canadensis</i>	Threatened	While no resident populations are known from Wisconsin, the species occasionally occurs in northern forested areas, and counties listed are those with the highest likelihood of occurrence.

	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Kirtland's warbler <i>Setophaga kirtlandii</i>	Endangered	Young jack pine stands (5 to 25 years old)
	Fassett's locoweed <i>Oxytropis campestris</i> var. <i>chartacea</i>	Threatened	Open sandy lakeshores
Brown	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Rufa red knot <i>(Calidris canutus rufa)</i>	Proposed Threatened	Along Green Bay
	Dwarf lake iris <i>Iris lacustris</i>	Threatened	Partially shaded sandy-gravelly soils on lakeshores
Buffalo	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Eastern massasauga <i>Sistrurus catenatus</i>	Candidate	Open to forested wetlands and adjacent uplands
	Higgins eye pearly mussel <i>(Lampsilis higginsii)</i>	Endangered	Mississippi River
	Sheepnose <i>(Plethobasus cyphus)</i>	Endangered	Mississippi River
	Spectaclecase <i>(Cumberlandia monodonta)</i>	Endangered	Mississippi River
Burnett	Canada lynx <i>Lynx canadensis</i>	Threatened	While no resident populations are known from Wisconsin, the species occasionally occurs in northern forested areas, and counties listed are those with the highest likelihood of occurrence.
	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Whooping crane <i>Grus americanus</i>	**Non-essential experimental population	Open wetlands and lakeshores

	Karner blue butterfly <i>Lycaeides melissa samuelis</i>	Endangered	Prairie, oak savanna, and jack pine areas with wild lupine
	Spectaclecase (<i>Cumberlandia monodonta</i>)	Endangered	St. Croix River
Calumet	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Whooping crane <i>Grus americanus</i>	**Non-essential experimental population	Open wetlands and lakeshores
Chippewa	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Spectaclecase (<i>Cumberlandia monodonta</i>)	Endangered	St. Croix River
	Karner blue butterfly <i>Lycaeides melissa samuelis</i>	Endangered	Prairie, oak savanna, and jack pine areas with wild lupine
Clark	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Whooping crane (<i>Grus americanus</i>)	**Non-essential experimental population	Open wetlands and lakeshores
	Karner blue butterfly <i>Lycaeides melissa samuelis</i>	Endangered	Prairie, oak savanna, and jack pine areas with wild lupine
Columbia	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Whooping crane (<i>Grus americanus</i>)	**Non-essential experimental population	Open wetlands and lakeshores
	Higgins eye pearly mussel (<i>Lampsilis higginsii</i>)	Endangered	Wisconsin River
	Sheepnose (<i>Plethobasus cyphus</i>)	Endangered	Wisconsin River

	Mead's milkweed (<i>Asclepias meadii</i>)	Threatened	Upland tallgrass prairie or glade/barren habitat Note: all the Mead's milkweed sites in Wisconsin are reintroduction attempts and occur on protected conservation lands.
	Prairie bush-clover (<i>Lespedeza leptostachya</i>)	Threatened	Dry to mesic prairies with gravelly soil
Crawford	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Whooping crane (<i>Grus americanus</i>)	**Non-essential experimental population	Open wetlands and lakeshores
	Eastern massasauga <i>Sistrurus catenatus</i>	Candidate	Open to forested wetlands and adjacent uplands
	Higgins eye pearly mussel (<i>Lampsilis higginsii</i>)	Endangered	Mississippi River
	Sheepnose (<i>Plethobasus cyphus</i>)	Endangered	Shallow areas in larger rivers and streams
	Spectaclecase (<i>Cumberlandia monodonta</i>)	Endangered	Mississippi River Note: EO for Crawford county is historic-last observation 1982
Dane	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Whooping crane (<i>Grus americanus</i>)	**Non-essential experimental population	Open wetlands and lakeshores
	Higgins eye pearly mussel (<i>Lampsilis higginsii</i>)	Endangered	Lower Wisconsin River
	Sheepnose (<i>Plethobasus cyphus</i>)	Endangered	Shallow areas in larger rivers and streams
	Eastern prairie fringed orchid (<i>Platanthera leucophaea</i>)	Threatened	Wet grasslands

	Mead's milkweed (<i>Asclepias meadii</i>)	Threatened	Upland tallgrass prairie or glade/barren habitat Note: all the Mead's milkweed sites in Wisconsin are reintroduction attempts and occur on protected conservation lands.
	Prairie bush-clover (<i>Lespedeza leptostachya</i>)	Threatened	Dry to mesic prairies with gravelly soil
Dodge	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Whooping crane (<i>Grus americanus</i>)	**Non-essential experimental population	Open wetlands and lakeshores
Door	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Hine's emerald dragonfly (<i>Somatochlora hineana</i>)	Endangered	Calcareous streams & associated wetlands overlying dolomite bedrock
	Hine's emerald dragonfly (<i>Somatochlora hineana</i>)	Critical Habitat	Critical Habitat Maps
	Pitcher's thistle <i>Cirsium pitcheri</i>	Threatened	Stabilized dunes, and blowout areas
	Dwarf lake iris <i>Iris lacustris</i>	Threatened	Partially shaded sandy-gravelly soils on lakeshores
Douglas	Canada lynx <i>Lynx canadensis</i>	Threatened	While no resident populations are known from Wisconsin, the species occasionally occurs in northern forested areas, and counties listed are those with the highest likelihood of occurrence.
	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Kirtland's warbler <i>Setophaga kirtlandii</i>	Endangered	Potential breeding in young jack pine stands (5 to 25 years old)
	Piping plover <i>Charadrius melodus</i>	Endangered	Sandy beaches; bare alluvial and dredge spoil islands
	Piping plover <i>Charadrius melodus</i>	Critical Habitat Designated	

	Rufa red knot (<i>Calidris canutus rufa</i>)	Proposed Threatened	Along Lake Superior
	Fassett's locoweed <i>Oxytropis campestris</i> var. <i>chartacea</i>	Threatened	Open sandy lakeshores
Dunn	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Sheepnose (<i>Plethobasus cyphus</i>)	Endangered	Chippewa River
Eau Claire	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Sheepnose (<i>Plethobasus cyphus</i>)	Endangered	Chippewa River
	Karner blue butterfly <i>Lycaeides melissa samuelis</i>	Endangered	Prairie, oak savanna, and jack pine areas with wild lupine
Florence	Canada lynx <i>Lynx canadensis</i>	Threatened	While no resident populations are known from Wisconsin, the species occasionally occurs in northern forested areas, and counties listed are those with the highest likelihood of occurrence.
	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
Fond du Lac	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Whooping crane (<i>Grus americanus</i>)	**Non-essential experimental population	Open wetlands and lakeshores
Forest	Canada lynx <i>Lynx canadensis</i>	Threatened	While no resident populations are known from Wisconsin, the species occasionally occurs in northern forested areas, and counties listed are those with the highest likelihood of occurrence.
	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.

Grant	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Whooping crane <i>Grus americanus</i>	**Non-essential experimental population	Open wetlands and lakeshores
	Higgins eye pearly mussel <i>(Lampsilis higginsii)</i>	Endangered	Lower Wisconsin and Mississippi Rivers
	Sheepnose <i>(Plethobasus cyphus)</i>	Endangered	Shallow areas in larger rivers and streams
	Spectaclecase <i>(Cumberlandia monodonta)</i>	Endangered	Mississippi River Note: EO for Grant county is historic-last observation 1982
	Hine's emerald dragonfly <i>(Somatochlora hineana)</i>	Endangered	Calcareous streams & associated wetlands overlying dolomite bedrock
	Mead's milkweed <i>(Asclepias meadii)</i>	Threatened	Upland tallgrass prairie or glade/barren habitat Note: all the Mead's milkweed sites in Wisconsin are reintroduction attempts and occur on protected conservation lands.
	Northern monkshood <i>(Aconitum noveboracense)</i>	Threatened	North facing slopes
	Prairie bush-clover <i>(Lespedeza leptostachya)</i>	Threatened	Dry to mesic prairies, with gravelly soil
Green	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Whooping crane <i>(Grus americanus)</i>	**Non-essential experimental population	Open wetlands and lakeshores
	Mead's milkweed <i>(Asclepias meadii)</i>	Threatened	Upland tallgrass prairie or glade/barren habitat Note: all the Mead's milkweed sites in Wisconsin are reintroduction attempts and occur on protected conservation lands.
	Prairie bush-clover <i>(Lespedeza leptostachya)</i>	Threatened	Dry to mesic prairies, with gravelly soil

Green Lake	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Whooping crane <i>Grus americanus</i>	**Non-essential experimental population	Open wetlands and lakeshores
	Karner blue butterfly <i>Lycaeides melissa samuelis</i>	Endangered	Prairie, oak savanna, and jack pine areas with wild lupine
	Poweshiek skipperling (<i>Oarisma poweshiek</i>)	Proposed Endangered and Proposed Critical Habitat Critical Habitat Maps	Native prairie
	Eastern prairie fringed orchid (<i>Platanthera leucophaea</i>)	Threatened	Wet grasslands
Iowa	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Whooping crane (<i>Grus americanus</i>)	**Non-essential experimental population	Open wetlands and lakeshores
	Higgins eye pearly mussel (<i>Lampsilis higginsii</i>)	Endangered	Lower Wisconsin Rivers
	Sheepnose Mussel (<i>Plethobasus cyphus</i>)	Endangered	Wisconsin River
	Hine's emerald dragonfly (<i>Somatochlora hineana</i>)	Endangered	Calcareous streams & associated wetlands overlying dolomite bedrock
	Mead's milkweed (<i>Asclepias meadii</i>)	Threatened	Upland tallgrass prairie or glade/barren habitat Note: all the Mead's milkweed sites in Wisconsin are reintroduction attempts and occur on protected conservation lands.
	Prairie bush-clover (<i>Lespedeza leptostachya</i>)	Threatened	Dry to mesic prairies with gravelly soil
Iron	Canada lynx <i>Lynx canadensis</i>	Threatened	While no resident populations are known from Wisconsin, the species occasionally occurs in northern forested areas, and counties listed are those with the highest likelihood of occurrence.

	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
Jackson	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Whooping crane Grus americanus	**Non-essential experimental population	Open wetlands and lakeshores
	Kirtland's warbler <i>Setophaga kirtlandii</i>	Endangered	Potential breeding in young jack pine stands (5 to 25 years old)
	Eastern massasauga Sistrurus catenatus	Candidate	Open to forested wetlands and adjacent uplands
	Karner blue butterfly <i>Lycaeides melissa samuelis</i>	Endangered	Prairie, oak savanna, and jack pine areas with wild lupine
Jefferson	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Whooping crane Grus americanus	**Non-essential experimental population	Open wetlands and lakeshores
	Eastern prairie fringed orchid (<i>Platanthera leucophaea</i>)	Threatened	Wet grasslands
Juneau	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Whooping crane Grus americanus	**Non-essential experimental population	Open wetlands and lakeshores Whooping cranes have nested in this county
	Eastern massasauga Sistrurus catenatus	Candidate	Open to forested wetlands and adjacent uplands
	Karner blue butterfly <i>Lycaeides melissa samuelis</i>	Endangered	Prairie, oak savanna, and jack pine areas with wild lupine
Kenosha	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.

	Whooping crane Grus americanus	**Non-essential experimental population	Open wetlands and lakeshores
	Eastern prairie fringed orchid (Platanthera leucophaea)	Threatened	Wet grasslands
Kewaunee	Northern long-eared bat Myotis septentrionalis	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Hine's emerald dragonfly (Somatochlora hineana)	Endangered	Calcareous streams & associated wetlands overlying dolomite bedrock
LaCrosse	Northern long-eared bat Myotis septentrionalis	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Whooping crane Grus americanus	**Non-essential experimental population	Open wetlands and lakeshores
	Eastern massasauga Sistrurus catenatus	Candidate	Open to forested wetlands and adjacent uplands
	Higgins eye pearly mussel (Lampsilis higginsii)	Endangered	Mississippi River
	Sheepnose (Plethobasus cyphus)	Endangered	Mississippi River
Lafayette	Northern long-eared bat Myotis septentrionalis	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Whooping crane Grus americanus	**Non-essential experimental population	Open wetlands and lakeshores
	Prairie bush-clover (Lespedeza leptostachya)	Threatened	Dry to mesic prairies with gravelly soil
Langlade	Northern long-eared bat Myotis septentrionalis	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
Lincoln	Northern long-eared bat Myotis septentrionalis	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.

Manitowoc	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Rufa red knot (<i>Calidris canutus rufa</i>)	Proposed Threatened	Along Lake Michigan
	Pitcher's <i>Cirsium pitcheri</i>	Threatened	Stabilized dunes, and blowout areas
	Piping plover <i>Charadrius melodus</i>	Endangered	Sandy beaches; bare alluvial and dredge spoil islands
	Piping plover <i>Charadrius melodus</i>	Critical Habitat Designated	
Marathon	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Whooping crane <i>Grus americanus</i>	**Non-essential experimental population	Open wetlands and lakeshores
Marinette	Canada lynx <i>Lynx canadensis</i>	Threatened	While no resident populations are known from Wisconsin, the species occasionally occurs in northern forested areas, and counties listed are those with the highest likelihood of occurrence.
	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Kirtland's warbler <i>Setophaga kirtlandii</i>	Endangered	Potential breeding in young jack pine stands (5 to 25 years old)
	Piping plover <i>Charadrius melodus</i>	Endangered	Sandy beaches; bare alluvial and dredge spoil islands
	Piping plover <i>Charadrius melodus</i>	Critical Habitat Designated	
Marquette	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Whooping crane <i>Grus americanus</i>	**Non-essential experimental population	Open wetlands and lakeshores
	Karner blue butterfly <i>Lycaeides melissa samuelis</i>	Endangered	Prairie, oak savanna, and jack pine areas with wild lupine

Menominee	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Karner blue butterfly <i>Lycaeides melissa samuelis</i>	Endangered	Prairie, oak savanna, and jack pine areas with wild lupine
Milwaukee	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Rufa red knot (<i>Calidris canutus rufa</i>)	Proposed Threatened	Along Lake Michigan
Monroe	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Whooping crane <i>Grus americanus</i>	**Non-essential experimental population	Open wetlands and lakeshores Whooping cranes have nested in this county
	Eastern massasauga <i>Sistrurus catenatus</i>	Candidate	Open to forested wetlands and adjacent uplands
	Karner blue butterfly <i>Lycaeides melissa samuelis</i>	Endangered	Prairie, oak savanna, and jack pine areas with wild lupine
	Northern monkshood <i>Aconitum noveboracense</i>	Threatened	North facing slopes
Oconto	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Rufa red knot (<i>Calidris canutus rufa</i>)	Proposed Threatened	Along Green Bay
	Whooping crane <i>Grus americanus</i>	**Non-essential experimental population	Open wetlands and lakeshores
	Karner blue butterfly <i>Lycaeides melissa samuelis</i>	Endangered	Prairie, oak savanna, and jack pine areas with wild lupine
Oneida	Canada lynx <i>Lynx canadensis</i>	Threatened	While no resident populations are known from Wisconsin, the species occasionally occurs in northern forested areas, and counties listed are those with the highest likelihood of occurrence.

	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
Outagamie	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Snuffbox <i>Epioblasma triquetra</i>	Endangered	Small to medium-sized creeks and some larger rivers, in areas with a swift current
Ozaukee	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Rufa red knot <i>(Calidris canutus rufa)</i>	Proposed Threatened	Along Lake Michigan
	Hine's emerald dragonfly <i>Somatochlora hineana</i>	Endangered	Calcareous streams & associated wetlands overlying dolomite bedrock
	Hine's emerald dragonfly <i>Somatochlora hineana</i>	Critical Habitat	Critical Habitat Maps
	Eastern prairie fringed orchid <i>Platanthera leucophaea</i>	Threatened	Wet grasslands
Pepin	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Whooping crane <i>Grus americanus</i>	**Non-essential experimental population	Open wetlands and lakeshores
	Eastern massasauga <i>Sistrurus catenatus</i>	Candidate	Open to forested wetlands and adjacent uplands
	Sheepnose <i>Plethobasus cyphus</i>	Endangered	Mississippi River
	Prairie bush-clover <i>Lespedeza leptostachya</i>	Threatened	Dry to mesic prairies with gravelly soil
Pierce	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Higgins eye pearly mussel <i>Lampsilis Higginsi</i>	Endangered	Mississippi and St. Croix Rivers

	Snuffbox <i>Epioblasma triquetra</i>	Endangered	Small to medium-sized creeks and some larger rivers, in areas with a swift current
	Spectaclecase <i>Cumberlandia monodonta</i>	Endangered	Large rivers
	Prairie bush-clover <i>Lespedeza leptostachya</i>	Threatened	Dry to mesic prairies with gravelly soil
Polk	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Whooping crane <i>Grus americanus</i>	**Non-essential experimental population	Open wetlands and lakeshores
	Higgins eye pearly mussel <i>Lampsilis higginsii</i>	Endangered	St. Croix Rivers
	Snuffbox <i>Epioblasma triquetra</i>	Endangered	Small to medium-sized creeks and some larger rivers, in areas with a swift current
	Spectaclecase <i>Cumberlandia monodonta</i>	Endangered	St. Croix River
	Winged mapleleaf <i>Quadrula fragosa</i>	Endangered	St. Croix River
Portage	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Karner blue butterfly <i>Lycaeides melissa samuelis</i>	Endangered	Prairie, oak savanna, and jack pine areas with wild lupine
	Fassett's locoweed <i>Oxytropis campestris</i> var. <i>chartacea</i>	Threatened	Open sandy lakeshores
Price	Canada lynx <i>Lynx canadensis</i>	Threatened	While no resident populations are known from Wisconsin, the species occasionally occurs in northern forested areas, and counties listed are those with the highest likelihood of occurrence.
	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
Racine	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.

	Rufa red knot <i>(Calidris canutus rufa)</i>	Proposed Threatened	Along Lake Michigan
Richland	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Whooping crane <i>Grus americanus</i>	**Non-essential experimental population	Open wetlands and lakeshores
	Higgins eye pearly mussel <i>Lampsilis higginsii</i>	Endangered	Lower Wisconsin River
	Sheepnose <i>Plethobasus cyphus</i>	Endangered	Wisconsin River
	Hine's emerald dragonfly <i>Somatochlora hineana</i>	Endangered	Calcareous streams & associated wetlands overlying dolomite bedrock
	Northern monkshood <i>Aconitum noveboracense</i>	Threatened	North facing slopes
Rock	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Whooping crane <i>Grus americanus</i>	**Non-essential experimental population	Open wetlands and lakeshores
	Eastern massasauga <i>Sistrurus catenatus</i>	Candidate	Open to forested wetlands and adjacent uplands
	Eastern prairie fringed orchid <i>Platanthera leucophaea</i>	Threatened	Wet grasslands
	Prairie bush-clover <i>Lespedeza leptostachya</i>	Threatened	Dry to mesic prairies with gravelly soil
Rusk	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Sheepnose <i>Plethobasus cyphus</i>	Endangered	Chippewa River
St. Croix	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.

	Higgins eye pearly mussel <i>Lampsilis higginsii</i>	Endangered	St. Croix River
	Snuffbox <i>Epioblasma triquetra</i>	Endangered	Small to medium-sized creeks and some larger rivers, in areas with a swift current
	Spectaclecase <i>Cumberlandia monodonta</i>	Endangered	St. Croix River
	Winged mapleleaf <i>Quadrula fragosa</i>	Endangered	St. Croix River
	Prairie bush-clover <i>Lespedeza leptostachya</i>	Threatened	Dry to mesic prairies with gravelly soil
Sauk	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Whooping crane <i>Grus americanus</i>	**Non-essential experimental population	Open wetlands and lakeshores
	Higgins eye pearly mussel <i>Lampsilis higginsii</i>	Endangered	Wisconsin River
	Sheepnose <i>Plethobasus cyphus</i>	Endangered	Wisconsin River
	Northern monkshood <i>Aconitum noveboracense</i>	Threatened	North facing slopes
	Prairie bush-clover <i>Lespedeza leptostachya</i>	Threatened	Dry to mesic prairies with gravelly soil
Sawyer	Canada lynx <i>Lynx canadensis</i>	Threatened	While no resident populations are known from Wisconsin, the species occasionally occurs in northern forested areas, and counties listed are those with the highest likelihood of occurrence.
	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
Shawano	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Whooping crane <i>Grus americanus</i>	**Non-essential experimental population	Open wetlands and lakeshores
	Snuffbox <i>Epioblasma triquetra</i>	Endangered	Small to medium-sized creeks and some larger rivers, in areas with a swift current

	Karner blue butterfly <i>Lycaeides melissa samuelis</i>	Endangered	Prairie, oak savanna, and jack pine areas with wild lupine
Sheboygan	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Pitcher's <i>Cirsium pitcheri</i>	Threatened	Stabilized dunes, and blowout areas
Taylor	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
Trempealeau	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Whooping crane <i>Grus americanus</i>	**Non-essential experimental population	Open wetlands and lakeshores
	Eastern massasauga <i>Sistrurus catenatus</i>	Candidate	Open to forested wetlands and adjacent uplands
	Higgins eye pearly mussel <i>Lampsilis higginsii</i>	Endangered	Mississippi River
Vernon	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Higgins eye pearly mussel <i>Lampsilis higginsii</i>	Endangered	Mississippi River
	Northern monkshood <i>Aconitum noveboracense</i>	Threatened	North facing slopes
Vilas	Canada lynx <i>Lynx canadensis</i>	Threatened	While no resident populations are known from Wisconsin, the species occasionally occurs in northern forested areas, and counties listed are those with the highest likelihood of occurrence.
	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Kirtland's warbler <i>Setophaga kirtlandii</i>	Endangered	Potential breeding in young jack pine stands (5 to 25 years old)

Walworth	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Whooping crane <i>Grus americanus</i>	**Non-essential experimental population	Open wetlands and lakeshores
	Eastern prairie fringed orchid <i>Platanthera leucophaea</i>	Threatened	Wet grasslands
	Eastern massasauga <i>Sistrurus catenatus</i>	Candidate	Open to forested wetlands and adjacent uplands
Washburn	Canada lynx <i>Lynx canadensis</i>	Threatened	While no resident populations are known from Wisconsin, the species occasionally occurs in northern forested areas, and counties listed are those with the highest likelihood of occurrence.
	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Kirtland's warbler <i>Setophaga kirtlandii</i>	Endangered	Potential breeding in young jack pine stands (5 to 25 years old)
Washington	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Whooping crane <i>Grus americanus</i>	**Non-essential experimental population	Open wetlands and lakeshores
Waukesha	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Poweshiek skipperling <i>Oarisma poweshiek</i>	Proposed Endangered and Proposed Critical Habitat Critical Habitat Maps	Native prairie
	Eastern prairie fringed orchid <i>Platanthera leucophaea</i>	Threatened	Wet grasslands

Waupaca	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Snuffbox <i>Epioblasma triquetra</i>	Endangered	Small to medium-sized creeks and some larger rivers, in areas with a swift current
	Karner blue butterfly <i>Lycaeides melissa samuelis</i>	Endangered	Prairie, oak savanna, and jack pine areas with wild lupine
Waushara	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Whooping crane <i>Grus americanus</i>	**Non-essential experimental population	Open wetlands and lakeshores
	Snuffbox <i>Epioblasma triquetra</i>	Endangered	Small to medium-sized creeks and some larger rivers, in areas with a swift current
	Karner blue butterfly <i>Lycaeides melissa samuelis</i>	Endangered	Prairie, oak savanna, and jack pine areas with wild lupine
	Fassett's locoweed <i>Oxytropis campestris</i> var. <i>chartacea</i>	Threatened	Open sandy lakeshores
Winnebago	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Whooping crane <i>Grus americanus</i>	**Non-essential experimental population	Open wetlands and lakeshores
	Eastern prairie fringed orchid <i>Platanthera leucophaea</i>	Threatened	Wet grasslands
Wood	Northern long-eared bat <i>Myotis septentrionalis</i>	Proposed as Endangered	Hibernates in caves and mines - swarming in surrounding wooded areas in autumn. During summer, roosts and forages in upland forests.
	Whooping crane <i>Grus americanus</i>	**Non-essential experimental population	wetlands and lakeshores Whooping cranes have nested in this county
	Eastern massasauga <i>Sistrurus catenatus</i>	Candidate	Open to forested wetlands and adjacent uplands
	Karner blue butterfly <i>Lycaeides melissa samuelis</i>	Endangered	Prairie, oak savanna, and jack pine areas with wild lupine

